

Monitoring Well Network Project Peace Region, British Columbia, Canada		EERI-4													
Energy and Environment Research Initiative Dept of Earth, Oceans, & Atmospheric Sciences University of British Columbia		Date drilled : 8/31/2018-9/01/2018 Location : Beryl Prairie, Kristoph Farm Equipment : Sonic drill (truck mounted) Logged by : Max Goetz Sampled by : Andrew Allen Drilled by : Mud Bay Drilling													
Depth in metres	Water info	Well1: 3" PVC, 0.020 slot Well2: 1" PVC, 0.020 slot Well3: 3/4" PE Elev.:	<table border="1"> <thead> <tr> <th colspan="4">GRAIN SIZE & LITHOLOGY</th> </tr> <tr> <th>gravel</th> <th>sand</th> <th>silt</th> <th>clay</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	GRAIN SIZE & LITHOLOGY				gravel	sand	silt	clay				
GRAIN SIZE & LITHOLOGY															
gravel	sand	silt	clay												
		DESCRIPTION	<table border="1"> <tr> <td> Top soil, black, plant roots present, 0% clasts Silty clay, dark brown clay with lesser light brown silt, rare clasts are granule size. <1% clasts Silt with clay interlayers, light brown silt with 2 cm interlayers of dark brown clay, some laminated silt observed. 0% clasts </td> </tr> <tr> <td> Silt to medium sand, coarsening sequence starting from silt, to very fine sand, to fine sand, to medium sand. Some interlayers of black organic present. 0% clasts </td> </tr> <tr> <td> Diamict, matrix coarsens from clayey silt to fine sand. Grey, dense, poorly sorted, clasts are mostly pebbles, rare cobbles. Clasts are often rounded with a flat side showing scour marks. 20-25% clasts </td> </tr> <tr> <td> Oxidized diamict, poorly sorted, dense coarse sandy-silt diamict with disappearance of high clast content. Orange coloured iron oxidation is potentially weathered in place of clasts. 0% clasts Sandy gravel, coarse to very coarse sand with angular pieces of pebble sized gravel. Clast-supported, matrix is very muddy, clay rich (might be from drilling water). Minor iron oxidation coating on gravel. 80% clasts. Oxidized diamict, same unit description as in oxidized diamict above. </td> </tr> <tr> <td> Siltstone, mostly dry, light grey, bedrock competency increases with depth. No bedding/lamination observed. </td> </tr> </table>	Top soil, black, plant roots present, 0% clasts Silty clay, dark brown clay with lesser light brown silt, rare clasts are granule size. <1% clasts Silt with clay interlayers, light brown silt with 2 cm interlayers of dark brown clay, some laminated silt observed. 0% clasts	Silt to medium sand, coarsening sequence starting from silt, to very fine sand, to fine sand, to medium sand. Some interlayers of black organic present. 0% clasts	Diamict, matrix coarsens from clayey silt to fine sand. Grey, dense, poorly sorted, clasts are mostly pebbles, rare cobbles. Clasts are often rounded with a flat side showing scour marks. 20-25% clasts	Oxidized diamict, poorly sorted, dense coarse sandy-silt diamict with disappearance of high clast content. Orange coloured iron oxidation is potentially weathered in place of clasts. 0% clasts Sandy gravel, coarse to very coarse sand with angular pieces of pebble sized gravel. Clast-supported, matrix is very muddy, clay rich (might be from drilling water). Minor iron oxidation coating on gravel. 80% clasts. Oxidized diamict, same unit description as in oxidized diamict above.	Siltstone, mostly dry, light grey, bedrock competency increases with depth. No bedding/lamination observed.							
Top soil, black, plant roots present, 0% clasts Silty clay, dark brown clay with lesser light brown silt, rare clasts are granule size. <1% clasts Silt with clay interlayers, light brown silt with 2 cm interlayers of dark brown clay, some laminated silt observed. 0% clasts															
Silt to medium sand, coarsening sequence starting from silt, to very fine sand, to fine sand, to medium sand. Some interlayers of black organic present. 0% clasts															
Diamict, matrix coarsens from clayey silt to fine sand. Grey, dense, poorly sorted, clasts are mostly pebbles, rare cobbles. Clasts are often rounded with a flat side showing scour marks. 20-25% clasts															
Oxidized diamict, poorly sorted, dense coarse sandy-silt diamict with disappearance of high clast content. Orange coloured iron oxidation is potentially weathered in place of clasts. 0% clasts Sandy gravel, coarse to very coarse sand with angular pieces of pebble sized gravel. Clast-supported, matrix is very muddy, clay rich (might be from drilling water). Minor iron oxidation coating on gravel. 80% clasts. Oxidized diamict, same unit description as in oxidized diamict above.															
Siltstone, mostly dry, light grey, bedrock competency increases with depth. No bedding/lamination observed.															

EERI-4